

Referral Criteria

NCCVEH at Prevent Blindness

- Ages 3 years:
 - *Majority of optotypes on 20/50 line*
- Ages 4 and 5 years:
 - *Majority of optotypes on 20/40 line*
- Ages 6 years and older:
 - *Majority of optotypes on 20/32 line*

AAP, AAPOS, AACO, AAO

- Ages 3 years:
 - *Majority of optotypes on 20/50 line*
- Ages 4 years:
 - *Majority of optotypes on 20/40 line*
- Ages 5 years and older:
 - *Majority of optotypes on 20/32 (or 20/30) line*
 - *2-line difference between eyes, even in passing area (i.e., 20/15 and 20/25) threshold screening*

Donahue, S. P., Baker, C. N., Committee on Practice and Ambulatory Medicine, Section on Ophthalmology, American Association of Certified Orthoptists, American Association for Pediatric Ophthalmology and Strabismus, American Academy of Ophthalmology (2016). Procedures for the evaluation of the visual system by pediatricians. *Pediatrics*, 137(1), 1-9. <https://pediatrics.aappublications.org/content/pediatrics/137/1/e20153597.full.pdf>

Cotter, S. A., Cyert, L. A., Miller, J. M., & Quinn, G. E. for the National Expert Panel to the National Center for Children's Vision and Eye Health. (2015). Vision screening for children 36 to <72 months: Recommended Practices. *Optometry and Vision Science*, 92(1), 6-16. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4274336/pdf/opx-92-06.pdf>